

Information for breastfeeding families



Calming a Colicky Breastfed Baby

Crying is often related to hunger, and feeding is the obvious solution. However, periods of crying with no obvious cause and no ready cure plague many parents. Here are a few suggestions. Give several a try. One may work today and another may work tomorrow. Be flexible and know that "this too will pass."

Symptoms:

Periods of inconsolable crying
Clenching fists
Drawing legs up to abdomen
Stiffening as if in pain
Passing gas
Grimacing

Possible causes: (may be more than one)

Immature gastro-intestinal system
Sensitive nervous system
Excessive gas or over-feeding
Formula intolerance
Foods the breastfeeding parent has eaten
Over-stimulation

Possible remedies:

- ✓ Hold your baby skin-to-skin several times each day
- ✓ Motion in any form
 - Walking
 - Baby swing
 - Rocking in rocking chair
 - Rocking side to side while standing
 - Ride in the car or in a stroller
 - Carry baby in infant sling
- ✓ Over-the-counter gas-relief drops (discuss with MD)
- ✓ Singing, humming
- ✓ Take a break, let someone else try for awhile; babies sense your tension
- ✓ White sound nearby (running water, vacuum, clothes dryer, hair dryer)
- ✓ Commercially available recordings with strong beat designed to simulate intrauterine sounds
- ✓ Undress your baby and allow complete freedom of motion
- ✓ Avoid over-stimulation from noises, lights, or motion
- ✓ Distract the baby with different sounds, sights, or places
- ✓ Place pressure on your baby's abdomen (gently)
 - Hold baby in the "colic hold" (facing floor, supported by your arm, heel of your hand putting pressure on the abdomen)
 - Hold baby over your shoulder or over your knees
- ✓ Bicycle your baby's legs
- ✓ Take an infant massage class
- ✓ Keep a food diary to determine if a particular food bothers your baby
- ✓ Consult a Lactation Consultant
- ✓ Contact your healthcare provider for other medical explanations and solutions

